



A cadavericfoetal study of craniorachichisis totalis

Latha M¹, Vino Victor Jesudas^{2*}, Froebel Giftly T.C³

¹Assistant Professor of Anatomy, ²Director and Professor of Anatomy and ³Tutor in Anatomy, Institute of Anatomy, Madurai Medical College, Madurai, Tamilnadu, India-625020

Received: 11-01-2018 / Revised Accepted: 30-01-2018 / Published: 01-02-2018


ABSTRACT

Aim of the study is to study in detail about Neural tube defects and the importance of folic acid during antenatal period. Neurulation is the process whereby the neural plate forms the neural tube. Cranial end is called anterior neuropore and caudal end is called Posterior neuropore. We received a male dead fetus from Institute of obstetrics and Gynaecology, Govt Rajaji Hospital Madurai. External examination of fetus showing Craniorachichisis was observed. Detailed study of fetus was performed. Craniorachichisis is a rare and most severe form of Neural tube defect in which both Brain and spinal cord remain open. Both Anencephaly and Spina bifida are present. Craniorachichisis can be diagnosed prenatally by ultrasound and by determination of Alpha fetoprotein (AFP) levels in maternal serum and amniotic fluid. It is estimated that 50-70% of Neural tube defects can be prevented if women take 400ug of Folic acid daily at least 1 month prior to conception and continuing throughout early pregnancy.

Keywords: Craniorachichisis, Ancencephaly, Spina Bifida, Neural tube defect (NTD)

Address for Correspondence: Dr.Vino Victor Jesudas, Director and Professor, Institute of Anatomy, Madurai Medical College, Madurai, Tamilnadu, India-625020; E-mail: vinovictor@rocketmail.com

How to Cite this Article: Latha M, Vino Victor Jesudas, Froebel Giftly T.C. A cadavericfoetal study of craniorachichisis totalis. World J Pharm Sci 2018; 6(2): 70-73.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which allows adapt, share and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms. 

© 2018 World J Pharm Sci

INTRODUCTION

Craniorachichisis is a rare and most severe form of Neural tube defect in which both Brain and spinal cord remain exposed .Both Anencephaly and spina bifida are present. Neural tube defects occur when the neural tube fail to close. If the neural tube fail to close in the cranial region, then most of the brain fail to develop .There is failure in the development of vault of the skull. This defect is called anencephaly. If the closure fails caudally anywhere from the cervical region to sacral region, then the defect is called Spina bifida.

Most common site for Spina bifida to occur is in the lumbosacral region. If the entire length of the neural tube remains open from the surface of the head on to the back, then the defect is called craniorachichisis totalis ^[1]. Miscarriage of the fetus commonly occurs during pregnancy .Usually fetus dies shortly after birth or stillbirth occurs. Some cases of craniorachichisis may be associated with other anomalies like Omphalocele, Congenital diaphragmatic hernia, Renal anomalies and Cardiac anomalies.^[2,3]

Origin of Neural tube defect is multifactorial and the likelihood of having a child with such defect increases significantly once one affected offspring is born. Folic acid deficiency^[1,6] plays an important role in occurrence of Neural tube defect. Incidence of Neural tube defects involving Brain or Spina cord or both is 0.5-2 per 1000 pregnancies worldwide. Incidence of neural tube defects in India ranges from 0.6-13/1000 live births and varies in different populations^[4]. More common in females than males. Polymorphism in the Methylenetetrahydrofolate reductase (MTHFR) gene can be a susceptibility factor in some

population. Reduction in MTHFR activity results in decreased plasma folate levels^[5] .Recently, mutations in the VANGL genes have been identified and associated with neural tube defects. Neural tube defects show an incidence of 5-10% in Trisomy 13 and Trisomy 18^[1]. Poor socioeconomic status, Multiparity, Maternal hyperthermia, Valproic acid exposure, Hypervitaminosis^[5]

MATERIALS AND METHODS

Of the ten fetuses received from Institute of Obstetrics and Gynaecology, Govt Rajaji Hospital Madurai during the year 2017 , one dead male fetus of 28weeks gestation having craniorachichisis totalis with anencephaly and spina bifida was observed. Academic autopsy was performed in Institute of Anatomy to see the presence of other associated congenital malformations. Their findings were appropriately documented.

OBSERVATION

The fetus on external examination showed craniorachichisis .It showed absence of scalp ,cranial vault. The defect extended upto lower thoracic region. Brain tissue and spinal cord in the cervical and lower thoracic region were exposed. Below the level of lower thoracic region no defect was observed and vertebral column was covered by normal skin. Eyes were bulging and there was absence of neck .Protruded tongue and prominent ears were observed. Fetus had extreme extension of head with Frog like appearance of face. Fetus had imperforate anus also .No other abnormality was found externally. Academic autopsy was carried out and no abnormalities were found internally.



Fig 1: Protruded tongue and absence of neck



Fig 2: Exposed Brain tissue and spinal cord



Fig 3: Prominent ears



Fig 4: Frog like appearance of face

DISCUSSION

Neurulation is the process whereby the neural plate forms the neural tube. By the end of the 3rd week, the lateral edges of the neural plate become elevated to form neural folds, and the midregion forms the neural groove. Gradually the neural folds approach each other in the midline, where they fuse. Fusion begins in the cervical region and proceeds cranially and caudally. As a result, the neural tube is formed. Cranial end is called anterior neuropore and caudal end is called posterior neuropore. Neural tube communicates with the neural cavity through ant and post neuropore. Closure of the cranial neuropore occurs at 25th day and posterior neuropore at 28th day. Hence CNS is represented by a closed tubular structure with a narrow caudal portion, the spinal cord and a broader cephalic portion characterized by a number of dilatations called Brain vesicles. Most defects of the spinal cord result from abnormal closure of neural folds in the 3rd and 4th weeks of development resulting in abnormalities involving meninges, vertebrae, muscles and skin. Recent evidence proved that Folic acid reduces the occurrence of Neural Tube Defects by as much as 50-70% if 400ug is taken daily atleast one month prior to conception and continued throughout early pregnancy.^[1,6]

Dr. Jose Ignacio (1992), reported a case of craniorachichisis associated with sirenomelia and hypoplasia of phlanges of right thumb. Ashutosh Halder, (2001) reported a case of sirenomelia associated with craniorachichisis. Associated cardiac abnormality (primitive heart), Renal agenesis, absent external genitalia were also observed. Kiran G (2009) reported a case of craniorachichisis totalis with only external abnormalities and they found no abnormalities internally. NS Naveen, (2010), observed a 13 weeks old female fetus with Craniorachichisis. sternal cleft and exomphalos were also seen with craniorachichisis, N.Ratnakar rao (2015)

reported a case of craniorachichisis with only external abnormalities.

Singh A, (2016) observed a 19 weeks old male fetus with congenital diaphragmatic hernia associated with craniorachichisis during 2017 we observed 10 fetuses ranging from 28-30 weeks. One fetus was observed with the following deformities, Frog like appearance of face, Protruded tongue, Protruding eyes with absence of neck, Absence of scalp and cranial vault, Deformed ears and Anal atresia. The incidence of craniorachichisis observed by us from our study was 0.5 %.

CONCLUSION

Craniorachichisis can be diagnosed prenatally by ultrasound and by determination of AFP levels in maternal serum and amniotic fluid^[6]. Vertebral column can be visualized by 12 weeks of gestation and the defect in closure of the vertebral arches can be detected. The prognosis of craniorachichisis is exceptionally poor and death of the neonate is unavoidable.

Experimental treatment is available only for spina bifida. It involves to perform surgery in uterus as early as 22 weeks gestation. The fetus is exposed by an incision into the uterus and the defect is repaired and infant is placed back in the uterus.^[1] In this case the mother belongs to low socioeconomic status and there was a history of non-consumption of Folic acid during antenatal period which could be the etiological factor for craniorachichisis. It's estimated that 50-70% of NTDs can be prevented if women take 400ug of Folic acid daily atleast 1 month prior to conception and continuing through early pregnancy. 50% of the pregnancies are unplanned so it is recommended that all women of child bearing age should take a multivitamin containing 400ug of folic acid daily.^[1,6]

REFERENCES

1. T.W.Saddler. Langman's Medical Embryology. Central nervous system, Neural tube defects, 13th Ed;Williams & Wilkins :Philadelphia,2012; 315-317.
2. Singh A et al. Craniorachischisis Totalis with Congenital Diaphragmatic Hernia. Fetal Paediatric Pathology.2016;35(3):192-8.
3. Ashutosh Halder et al. Sirenomelia associated with craniorachichisis totalis, Limb reduction and Primitive heart,Journal of Indian Paediatrics .2001;38:1041-1045.
4. Gayathri P et al .A fetal study of craniorachichisis with emphasis on Prenatal Diagnosis and Prevention.JMSCR.201; 04(4):10140-145.
5. Peter Turnpenny, Sian Ellard. Emery's Elements of Medical Genetics. Congenital abnormalities and dysmorphic syndrome,Neural tube defects,14th Ed;Churchill Livingstone:Edinburgh , 2007:258-260.
6. William J Larsen. Larsen's Human Embryology, Neural tube defects,5th Ed;Churchill Livingstone:Hongkong,2008:92-96.
7. Jaganmohan D et al. Two cases of craniospinal rachischisis totalis: Role of magnetic resonance imaging in diagnosis and review of neural tube defects in the Indian context with implications for folate fortification. J Pediatr Neurosci 2017;12:32-5.
8. Coskun A et al. Craniorachichisis Totalis-Case report and Review.Fetal Diagnosis and Theraphy.2009;25:21-25.
9. N.S. Naveen and Murlimanju. Craniorachischisis totalis.Journal of Neurosciences and Rural Practice .2010; 1(1): 54–55.
10. Rodríguez and Palacios J. Craniorachischisis totalis and sirenomelia. Am J Med Genet.1992;43(4):732-6.
11. Himabindu N et al. Comprehensive study of Neural Tube Defect in 1000 fetuses with clinical spectrum. Int J Anat Res 2015;3(4):1456-62